

Ground states for the pseudo-relativistic Hartree equation with external potential

Silvia Cingolani^{*}(s.cingolani@poliba.it)

Simone Secchi[†](simone.secchi@unimib.it)

(MS received ;)

We prove existence of positive ground state solutions to the pseudo-relativistic Schrödinger equation

$$\begin{cases} \sqrt{-\Delta + m^2}u + Vu = (W * |u|^\theta) |u|^{\theta-2}u & \text{in } \mathbb{R}^N \\ u \in H^{1/2}(\mathbb{R}^N) \end{cases}$$

where $N \geq 3$, $m > 0$, V is a bounded external scalar potential and W is a convolution potential, radially symmetric, satisfying suitable assumptions. We also furnish some asymptotic decay estimates of the found solutions.

1. Introduction

The mean field limit of a quantum system describing many self-gravitating, relativistic bosons with rest mass $m > 0$ leads to the time-dependent pseudo-relativistic Hartree equation

$$i \frac{\partial \psi}{\partial t} = \left(\sqrt{-\Delta + m^2} - m \right) \psi - \left(\frac{1}{|x|} * |\psi|^2 \right) \psi, \quad x \in \mathbb{R}^3 \quad (1.1)$$

where $\psi : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$ is the wave field. Such a physical system is often referred to as a boson star in astrophysics (see [12, 13, 14]). Solitary wave solutions $\psi(t, x) = e^{-it\lambda} \phi$, $\lambda \in \mathbb{R}$ to equation (1.1) satisfy the equation

$$\left(\sqrt{-\Delta + m^2} - m \right) \phi - \left(\frac{1}{|x|} * |\phi|^2 \right) \phi = \lambda \phi \quad (1.2)$$

For the non-relativistic Hartree equation, existence and uniqueness (modulo translations) of a minimizer has been proved by Lieb [17] by using symmetric decreasing rearrangement inequalities. Within the same setting, always for the negative Laplacian, Lions [21] proved existence of infinitely many spherically symmetric solutions by application of abstract critical point theory both without the constraint and with the constraint for a more general radially symmetric convolution potential. The non-relativistic Hartree equation is also known as the Choquard-Pekar or

^{*}Permanent address: Dipartimento di Meccanica, Matematica e Management, Politecnico di Bari. Via Orabona 4, 70125 Bari (Italy)

[†]Permanent address: Dipartimento di matematica e applicazioni, Università di Milano Bicocca. Via R. Cozzi 53, 20125 Milano (Italy)

Schrödinger-Newton equation and recently a large amount of papers are devoted to the study of solitary states and its semiclassical limit (see [1, 5, 7, 8, 9, 13, 18, 24, 22, 27, 28, 29, 31, 32] and references therein).

In [20] Lieb and Yau solved the pseudo-relativistic Hartree equation (1.2) by minimization on the sphere $\{\phi \in L^2(\mathbb{R}^3) \mid \int_{\mathbb{R}^3} |\phi|^2 = M\}$, and they proved that a radially symmetric ground state exists in $H^{1/2}(\mathbb{R}^3)$ whenever $M < M_c$, the so-called Chandrasekhar mass. These results have been generalized in [10]. Later Lenzmann proved in [16] that this ground state is unique (up to translations and phase change) provided that the mass M is sufficiently small; some results about the non-degeneracy of the ground state solution are also given.

Quite recently, Coti Zelati and Nolasco (see [11]) studied the equation

$$\sqrt{-\Delta + m^2}u = \mu u + \nu |u|^{p-2}u + \sigma(W * u^2)u$$

under the assumptions that $p \in \left(2, \frac{2N}{N-1}\right)$, $N \geq 3$, $\mu < m$, $m > 0$, $\nu \geq 0$, $\sigma \geq 0$ but not both zero, $W \in L^r(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$, $W \geq 0$, $r > N/2$, W is radially symmetric and decays to zero at infinity. They proved the existence of a positive, radial solution that decays to zero at infinity exponentially fast. For the case $\sigma < 0$, $\mu < m$, we also refer to [26] where a more general nonlinear term is considered.

In the present work we consider a generalized pseudo-relativistic Hartree equation

$$\sqrt{-\Delta + m^2}u + Vu = (W * |u|^\theta) |u|^{\theta-2}u \quad \text{in } \mathbb{R}^N \quad (1.3)$$

where $N \geq 3$, $m > 0$, V is an external potential and $W \geq 0$ is a radially symmetric convolution kernel such that $\lim_{|x| \rightarrow +\infty} W(|x|) = 0$.

In [23] Melgaard and Zongo prove that (1.3) has a sequence of radially symmetric solutions of higher and higher energy, assuming that V is radially symmetric potential, $\theta = 2$ and under some restrictive assumptions on the structure of the kernel W .

Here we are interested to look for positive ground state solutions for the pseudo-relativistic Hartree equation (1.3) when V is not symmetric. In such a case, the nonlocality of $\sqrt{-\Delta + m^2}$ and the presence of the external potential V (not symmetric) complicate the analysis of the pseudo-relativistic Hartree equation in a substantial way. The main difficulty is, as usual, the lack of compactness.

In what follows, we will keep the following assumptions:

(V1) $V: \mathbb{R}^N \rightarrow \mathbb{R}$ is a continuous and bounded function, and $V(y) + V_0 \geq 0$ for every $y \in \mathbb{R}^N$ and for some $V_0 \in (0, m)$.

(V2) There exist $R > 0$ and $k \in (0, 2m)$ such that

$$V(x) \leq V_\infty - e^{-k|x|} \quad \text{for all } |x| \geq R \quad (1.4)$$

where $V_\infty = \liminf_{|x| \rightarrow +\infty} V(x) > 0$.

(W) $W \in L^r(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$ for some $r > \max\left\{1, \frac{N}{N(2-\theta)+\theta}\right\}$ and $2 \leq \theta < \frac{2N}{N-1}$.

Our main result is the following.

Theorem 1.1. *Retain assumptions (V1), (V2) and (W). Then equation (1.3) has at least a positive solution $u \in H^{1/2}(\mathbb{R}^N)$.*

We remark that Theorem 1.1 applies for a large class of bounded electric potentials without symmetric constraints and covers the physically relevant cases of Newton or Yukawa type two body interaction, i.e. $W(x) = \frac{1}{|x|^\lambda}$ with $0 < \lambda < 2$, $W(x) = \frac{e^{-|x|}}{|x|}$.

The theorem will be proved using variational methods. Firstly we transform the problem into an elliptic equation with nonlinear Neumann boundary conditions, using a *local* realization of the pseudo-differential operator $\sqrt{-\Delta + m^2}$ as in [3, 4, 11, 10]. The corresponding solutions are found as critical points of an Euler functional defined in $H^1(\mathbb{R}_+^{N+1})$. We show that such a functional satisfies the Palais-Smale condition below some energy level determined by the value of V at infinity and we prove the existence of a Mountain Pass solution under the level where the Palais-Smale condition holds. Finally we also furnish some asymptotic decay estimates of the found solution.

Local and global well-posedness results for pseudo-relativistic Hartree equations with external potential are proved by Lenzmann in [15].

2. The variational framework

Before we state our main result, we recall a few basic facts about the functional setting of our problem. The operator $\sqrt{-\Delta + m^2}$ can be defined by Fourier analysis: given any $\phi \in L^2(\mathbb{R}^3)$ such that

$$\int_{\mathbb{R}^3} (m^2 + |k|^2) |\mathcal{F}\phi|^2 dk < +\infty,$$

we define $\sqrt{-\Delta + m^2}\phi$ via the identity

$$\mathcal{F} \left(\sqrt{-\Delta + m^2}\phi \right) = \sqrt{m^2 + |k|^2} \mathcal{F}\phi,$$

$\mathcal{F}$ being the usual Fourier transform. The condition

$$\int_{\mathbb{R}^3} \sqrt{m^2 + |k|^2} |\mathcal{F}\phi|^2 dk < +\infty$$

is known to be equivalent to $\phi \in H^{1/2}(\mathbb{R}^3)$. In this sense, the fractional Sobolev space $H^{1/2}(\mathbb{R}^3)$ is the natural space to work in. However, this definition is not particularly convenient for variational methods, and we prefer a *local* realization of the operator in the augmented half-space, originally inspired by the paper [6] for the fractional laplacian.

Given $u \in \mathcal{S}(\mathbb{R}^N)$, the Schwarz space of rapidly decaying smooth functions defined on $\mathbb{R}^N$, there exists one and only one function $v \in \mathcal{S}(\mathbb{R}_+^{N+1})$ (where $\mathbb{R}_+^{N+1} = (0, +\infty) \times \mathbb{R}^N$) such that

$$\begin{cases} -\Delta v + m^2 v = 0 & \text{in } \mathbb{R}_+^{N+1} \\ v(0, y) = u(y) & \text{for } y \in \mathbb{R}^N = \partial\mathbb{R}_+^{N+1}. \end{cases}$$

Setting

$$Tu(y) = -\frac{\partial v}{\partial x}(0, y),$$

we easily see that the problem

$$\begin{cases} -\Delta w + m^2 w = 0 & \text{in } \mathbb{R}_+^{N+1} \\ w(0, y) = Tu(y) & \text{for } y \in \partial\mathbb{R}_+^{N+1} = \{0\} \times \mathbb{R}^N \simeq \mathbb{R}^N \end{cases}$$

is solved by $w(x, y) = -\frac{\partial v}{\partial x}(x, y)$. From this we deduce that

$$T(Tu)(y) = -\frac{\partial w}{\partial x}(0, y) = \frac{\partial^2 v}{\partial x^2}(0, y) = (-\Delta_y v + m^2 v)(0, y),$$

and hence $T \circ T = (-\Delta_y + m^2)$, namely T is a square root of the Schrödinger operator $-\Delta + m^2$ on $\mathbb{R}^N = \partial\mathbb{R}_+^{N+1}$.

In the sequel, we will write $|\cdot|_p$ for the norm in $L^p(\mathbb{R}^N)$ and $\|\cdot\|_p$ for the norm in $L^p(\mathbb{R}_+^{N+1})$. The symbol $\|\cdot\|$ will be reserved for the usual norm of $H^1(\mathbb{R}_+^{N+1})$.

The theory of traces for Sobolev spaces ensures that every function $v \in H^1(\mathbb{R}_+^{N+1})$ possesses a *trace* $\gamma(v) \in H^{1/2}(\mathbb{R}^N)$ which satisfies the inequality (see [30, Lemma 13.1])

$$|\gamma(v)|_p^p \leq p \|v\|_{2(p-1)}^{p-1} \left\| \frac{\partial v}{\partial x} \right\|_2 \quad (2.1)$$

whenever $2 \leq p \leq 2N/(N-1)$. This also implies that, for every $\lambda > 0$,

$$\int_{\mathbb{R}^N} \gamma(v)^2 \leq \lambda \int_{\mathbb{R}_+^{N+1}} |v|^2 + \frac{1}{\lambda} \int_{\mathbb{R}_+^{N+1}} \left| \frac{\partial v}{\partial x} \right|^2. \quad (2.2)$$

As a particular case, we record

$$\int_{\mathbb{R}^N} \gamma(v)^2 \leq m \int_{\mathbb{R}_+^{N+1}} |v|^2 + \frac{1}{m} \int_{\mathbb{R}_+^{N+1}} |\nabla v|^2. \quad (2.3)$$

It is also known (see [30, Lemma 16.1]) that any element of $H^{1/2}(\mathbb{R}^N)$ is the trace of some function in $H^1(\mathbb{R}_+^{N+1})$.

From the previous construction and following [3, 4, 11, 10], we can replace the *nonlocal* problem (1.3) with the local Neumann problem

$$\begin{cases} -\Delta v + m^2 v = 0 & \text{in } \mathbb{R}_+^{N+1} \\ -\frac{\partial v}{\partial x} = -V(y)v + (W * |v|^\theta)|v|^{\theta-2}v & \text{in } \mathbb{R}^N = \partial\mathbb{R}_+^{N+1}. \end{cases} \quad (2.4)$$

We will look for solutions to (2.4) as critical points of the Euler functional $I: H^1(\mathbb{R}_+^{N+1}) \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} I(v) = \frac{1}{2} \iint_{\mathbb{R}_+^{N+1}} (|\nabla v|^2 + m^2 v^2) dx dy + \frac{1}{2} \int_{\mathbb{R}^N} V(y) \gamma(v)^2 dy \\ - \frac{1}{2\theta} \int_{\mathbb{R}^N} (W * |\gamma(v)|^\theta) |\gamma(v)|^\theta dy. \end{aligned} \quad (2.5)$$

We recall the well-known Young's inequality.

Proposition 2.1. Assume $f \in L^p(\mathbb{R}^N)$, $g \in L^q(\mathbb{R}^N)$ with $1 \leq p, q, r \leq \infty$, $p^{-1} + q^{-1} = 1 + r^{-1}$. Then

$$|f * g|_r \leq |f|_p |g|_q. \quad (2.6)$$

A major tool for our analysis is the generalized Hardy-Littlewood-Sobolev inequality. We recall that $L_w^q(\mathbb{R}^N)$ is the *weak* L^q space: see [19] for a definition. We denote by $|\cdot|_{q,w}$ the usual norm in $L_w^q(\mathbb{R}^N)$.

Proposition 2.2 ([19]). Assume that p, q and t lie in $(1, +\infty)$ and $p^{-1} + q^{-1} + t^{-1} = 2$. Then, for some constant $N_{p,q,t} > 0$ and for any $f \in L^p(\mathbb{R}^N)$, $g \in L^t(\mathbb{R}^N)$ and $h \in L_w^q(\mathbb{R}^N)$, we have the inequality

$$\left| \int f(x) h(x-y) g(y) dx dy \right| \leq N_{p,q,t} |f|_p |g|_t |h|_{q,w}. \quad (2.7)$$

For the sake of completeness, we check that the functional I is well-defined. As a consequence of (2.1), for every $p \in \left[2, \frac{2N}{N-1}\right]$ we deduce that

$$|\gamma(v)|_p \leq \frac{p-1}{p} \|v\|_{2(p-1)} + \|\nabla v\|_2 \leq C_p \|v\|, \quad (2.8)$$

and the term $\int_{\mathbb{R}^N} V(y) \gamma(v)^2 dy$ in the expression of $I(v)$ is finite because of the boundedness of V . Writing $W = W_1 + W_2 \in L^r(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$ and using (2.7) we can estimate the convolution term as follows:

$$\begin{aligned} \int_{\mathbb{R}^N} (W * |\gamma(v)|^\theta) |\gamma(v)|^\theta &= \int_{\mathbb{R}^N} (W_1 * |\gamma(v)|^\theta) |\gamma(v)|^\theta + \int_{\mathbb{R}^N} (W_2 * |\gamma(v)|^\theta) |\gamma(v)|^\theta \\ &\leq |W_1|_r |\gamma(v)|_{\frac{2r\theta}{2r-1}}^{2\theta} + |W_2|_\infty |\gamma(v)|_\theta^{2\theta} \leq |W_1|_r \|v\|^{2\theta} + |W_2|_\infty \|v\|^{2\theta}. \end{aligned} \quad (2.9)$$

Since

$$r > \frac{N}{N(2-\theta) + \theta} \quad \text{and} \quad 2 \leq \theta < \frac{2N}{N-1},$$

there results

$$\frac{2r-1}{2\theta r} = \frac{1}{\theta} - \frac{1}{2\theta r} > \frac{1}{\theta} - \frac{N(2-\theta) + \theta}{2\theta N} = \frac{N-1}{2N},$$

and thus

$$\frac{2\theta r}{2r-1} < \frac{2N}{N-1},$$

and from (2.9) we see that the convolution term in I is finite. It is easy to check, by the same token, that $I \in C^1(H^1(\mathbb{R}_+^{N+1}))$.

Remark 2.3. As we have just seen, estimates involving the kernel W always split into two parts. As a rule, those with the bounded kernel $W_2 \in L^\infty(\mathbb{R}^N)$ are straightforward. In the sequel, we will often focus on the contribution of $W_1 \in L^r(\mathbb{R}^N)$ and drop the easy computation with W_2 .

3. The limit problem

Let us consider the space of the symmetric functions

$$H^\sharp = \{u \in H^1(\mathbb{R}_+^{N+1}) \mid u(x, Ry) = u(x, y) \text{ for all } R \in O(N)\}.$$

Let us consider the functional $J_\alpha: H^\sharp \rightarrow \mathbb{R}$ defined by setting

$$J_\alpha(v) = \frac{1}{2} \iint_{\mathbb{R}_+^{N+1}} (|\nabla v|^2 + m^2 v^2) dx dy + \frac{1}{2} \int_{\mathbb{R}^N} \alpha \gamma(v)^2 dy - \frac{1}{2\theta} \int_{\mathbb{R}^N} (W * |\gamma(v)|^\theta) |\gamma(v)|^\theta dy. \quad (3.1)$$

where $W \geq 0$ is radially symmetric, $\lim_{|x| \rightarrow +\infty} W(|x|) = 0$ and assumption (W) holds. If $\alpha > -m$, we can extend the arguments in Theorem 4.3 [11], for the case $\theta = 2$, and prove that the functional J_α has a Mountain Pass critical point $v_\alpha \in H^\sharp$, namely

$$J_\alpha(v_\alpha) = E_\alpha = \inf_{g \in \Gamma_\sharp} \max_{t \in [0,1]} J_\alpha(g(t)) \quad (3.2)$$

where $\Gamma_\sharp = \{g \in C([0,1]; H^\sharp) \mid g(0) = 0, J_\alpha(g(1)) < 0\}$. The critical point v_α corresponds to a weak solution of

$$\begin{cases} -\Delta v + m^2 v = 0 & \text{in } \mathbb{R}_+^{N+1} \\ -\frac{\partial v}{\partial x} = -\alpha v + (W * |v|^\theta) |v|^{\theta-2} v & \text{in } \mathbb{R}^N = \partial \mathbb{R}_+^{N+1}. \end{cases} \quad (3.3)$$

In the sequel, we need a standard characterization of the mountain-pass level E_α . Let us define the *Nehari manifold* $\mathcal{N}_\alpha$ associated to the functional J_α :

$$\begin{aligned} \mathcal{N}_\alpha &= \left\{ v \in H^\sharp \mid \iint_{\mathbb{R}^N \times \mathbb{R}^N} |\nabla v|^2 + m^2 v^2 dx dy \right. \\ &\quad \left. = -\alpha \int_{\mathbb{R}^N} \gamma(v)^2 dy + \int_{\mathbb{R}^N} (W * |\gamma(v)|^\theta) |\gamma(v)|^\theta dy \right\}. \end{aligned} \quad (3.4)$$

Lemma 3.1. *There results*

$$\inf_{v \in \mathcal{N}_\alpha} J_\alpha(v) = \inf_{v \in H^\sharp} \max_{t > 0} J_\alpha(tv) = E_\alpha. \quad (3.5)$$

Proof. The proof is straightforward, since J_α is the sum of homogeneous terms; we follow [33]. First of all, for $v \in H^\sharp$ we compute

$$\begin{aligned} J_\alpha(tv) &= \frac{t^2}{2} \left(\iint_{\mathbb{R}^N \times \mathbb{R}^N} |\nabla v|^2 + m^2 v^2 dx dy + \alpha \int_{\mathbb{R}^N} \gamma(v)^2 dy \right) \\ &\quad - \frac{t^{2\theta}}{2\theta} \int_{\mathbb{R}^N} (W * |\gamma(v)|^\theta) |\gamma(v)|^\theta dy. \end{aligned} \quad (3.6)$$

Since $\theta \geq 2$ and by using (2.3), it is easy to check that $t \in (0, +\infty) \mapsto J(tv)$ possesses a unique critical point $t = t(v) > 0$ such that $t(v)v \in \mathcal{N}_\alpha$. Moreover, since J_α has the mountain-pass geometry, $t = t(v)$ is a maximum point. It follows that

$$\inf_{v \in \mathcal{N}_\alpha} J_\alpha(v) = \inf_{v \in H^\sharp} \max_{t > 0} J_\alpha(tv).$$

The manifold $\mathcal{N}_\alpha$ splits $H^\sharp$ into two connected components, and the component containing 0 is open. In addition, J_α is non-negative on this component, because $\langle J'_\alpha(tv), v \rangle \geq 0$ when $0 < t \leq t(v)$. It follows immediately that any path $\gamma: [0, 1] \rightarrow H^\sharp$ with $\gamma(0) = 0$ and $J_\alpha(\gamma(1)) < 0$ must cross $\mathcal{N}_\alpha$, so that

$$E_\alpha \geq \inf_{v \in \mathcal{N}_\alpha} J_\alpha(v).$$

The proof of (3.5) is complete. $\square$

Following completely analogous argument in Theorem 3.14 and Theorem 5.1 in [11], we can state the following result.

Theorem 3.2. *Let $\alpha + m > 0$ and (W) holds. Then $v_\alpha \in C^\infty([0, +\infty) \times \mathbb{R}^N)$, $v_\alpha(x, y) > 0$ in $[0, \infty) \times \mathbb{R}^N$ and for any $0 \leq \sigma \in (-\alpha, m)$ there exists $C > 0$ such that*

$$0 < v_\alpha(x, y) \leq C e^{-(m-\sigma)\sqrt{x^2+y^2}} e^{-\sigma x}$$

for all $(x, y) \in [0, +\infty) \times \mathbb{R}^N$. In particular,

$$0 < v_\alpha(0, y) \leq C e^{-\delta|y|} \quad \text{for every } y \in \mathbb{R}^N$$

where $0 < \delta < m + \alpha$ if $\alpha \leq 0$, and $\delta = m$ if $\alpha > 0$.

4. The Palais-Smale condition

For any $v \in H^1(\mathbb{R}_+^{N+1})$ we denote

$$\mathbb{D}(v) = \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) |\gamma(v)(x)|^\theta |\gamma(v)(y)|^\theta dx dy.$$

Inequality (2.9) yields immediately

$$\mathbb{D}(v) \leq K \|v\|^{2\theta} \quad (4.1)$$

for every $v \in H^1(\mathbb{R}_+^{N+1})$.

Lemma 4.1. *Let $\{v_n\}_n$ be a sequence in $H^1(\mathbb{R}_+^{N+1})$ such that $v_n \rightharpoonup 0$ weakly in $H^1(\mathbb{R}_+^{N+1})$*

$$I(v_n) \rightarrow c < E_{V_\infty} \quad \text{and} \quad I'(v_n) \rightarrow 0$$

where $V_\infty := \liminf_{|x| \rightarrow \infty} V(x) > 0$. Then a subsequence of $\{v_n\}_n$ converges strongly to 0 in $H^1(\mathbb{R}_+^{N+1})$.

Proof. First of all, we recall (2.3), and we rewrite $I(v)$ as

$$I(v) = \frac{1}{2} \int_{\mathbb{R}_+^{N+1}} |\nabla v|^2 + m^2 |v|^2 + \frac{1}{2} \int_{\mathbb{R}^N} (V + V_0) \gamma(v)^2 - \frac{V_0}{2} \int_{\mathbb{R}^N} \gamma(v)^2 - \frac{1}{2\theta} \mathbb{D}(v),$$

so that $V + V_0 \geq 0$ everywhere. Now,

$$c + 1 + \|v_n\| \geq I(v_n) - \frac{1}{2} \langle I'(v_n), v_n \rangle = \left(\frac{1}{2} - \frac{1}{2\theta} \right) \mathbb{D}(v_n), \quad (4.2)$$

which implies that, for some constants C_1 and C_2 ,

$$\frac{1}{2\theta} \mathbb{D}(v_n) \leq C_1 \|v_n\| + C_2.$$

But then, using (2.3),

$$c + 1 \geq I(v_n)$$

$$\begin{aligned} &\geq \frac{1}{2} \int_{\mathbb{R}_+^{N+1}} |\nabla v_n|^2 + \frac{m^2}{2} \int_{\mathbb{R}_+^{N+1}} |v_n|^2 - \frac{V_0}{2} \left(m \int_{\mathbb{R}_+^{N+1}} |v_n|^2 + \frac{1}{m} \int_{\mathbb{R}_+^{N+1}} |\nabla v_n|^2 \right) \\ &\quad - C_1 \|v_n\| - C_2 \\ &= \frac{1}{2} \left(1 - \frac{V_0}{m} \right) \int_{\mathbb{R}_+^{N+1}} |\nabla v_n|^2 + \frac{m(m - V_0)}{2} \int_{\mathbb{R}_+^{N+1}} |v_n|^2 - C_1 \|v_n\| - C_2 \end{aligned}$$

and since $m - V_0 > 0$ we deduce that $\{v_n\}$ is a bounded sequence in $H^1(\mathbb{R}_+^{N+1})$.

A standard argument shows that $\|v_n\|$ is bounded in $H^1(\mathbb{R}_+^{N+1})$ and

$$\frac{\theta - 1}{2\theta} \left(\|v_n\|^2 + \int_{\mathbb{R}^N} V(y) \gamma(v_n)^2 dy \right) \rightarrow c \quad \text{and} \quad \frac{\theta - 1}{2\theta} \mathbb{D}(v_n) \rightarrow c.$$

Therefore $c \geq 0$. If $c = 0$, then

$$\begin{aligned} o(1) &= \left(\|v_n\|^2 + \int_{\mathbb{R}^N} V(y) \gamma(v_n)^2 dy \right) \\ &\geq \left(1 - \frac{V_0}{m} \right) \int_{\mathbb{R}_+^{N+1}} |\nabla v_n|^2 + m(m - V_0) \int_{\mathbb{R}_+^{N+1}} |v_n|^2, \end{aligned}$$

and $m - V_0 > 0$ yields that $v_n \rightarrow 0$ strongly in $H^1(\mathbb{R}_+^{N+1})$.

Assume therefore that $c > 0$. Fix $\alpha < V_\infty$ such that $c < E_\alpha$, and $R_0 > 0$ such that $V(x) \geq \alpha$ if $|x| \geq R_0$. Let $\varepsilon \in (0, 1)$. Since $\{v_n\}_n$ is bounded in $H^1(\mathbb{R}_+^{N+1})$ there exists $R_\varepsilon > R_0$ such that $R_\varepsilon \rightarrow +\infty$ as $\varepsilon \rightarrow 0$ and, after passing to a subsequence,

$$\iint_{S_{R_\varepsilon}} (|\nabla v_n|^2 + m^2 v_n^2) dx dy + \int_{A_{R_\varepsilon}} V(y) \gamma(v_n)^2 dy < \varepsilon \quad \text{for all } n \in \mathbb{N}. \quad (4.3)$$

where

$$\begin{aligned} S_{R_\varepsilon} &= \{z = (x, y) \in \mathbb{R}_+^{N+1} \mid R_\varepsilon < |z| < R_\varepsilon + 1\} \\ A_{R_\varepsilon} &= \{y \in \mathbb{R}^N \mid R_\varepsilon < |y| < R_\varepsilon + 1\}. \end{aligned}$$

If this is not the case, for any $m \in \mathbb{N}$, $m \geq R_0$ there exists $\nu(m) \in \mathbb{N}$ such that

$$\iint_{S_m} (|\nabla v_n|^2 + m^2 v_n^2) dx dy + \int_{A_m} V(y) \gamma(v_n)^2 dy \geq \varepsilon \quad (4.4)$$

for any $n \in \mathbb{N}$, $n \geq \nu(m)$. We can assume that $\nu(m)$ is nondecreasing. Therefore for any integer $m \geq R_0$ there exists an integer $\nu(m)$ such that

$$\begin{aligned} \|v_n\|^2 + \int_{\mathbb{R}^N} V(y) \gamma(v_n)^2 dy &\geq \iint_{T_m} (|\nabla v_n|^2 + m^2 v_n^2) dx dy + \int_{B_m} V(y) \gamma(v_n)^2 dy \\ &\geq (m - R_0) \varepsilon \quad (4.5) \end{aligned}$$

for any $n \geq \nu(m)$, where $T_m = \{z = (x, y) \in \mathbb{R}_+^{N+1} \mid R_0 < |z| < m\}$ and $B_m = \{y \in \mathbb{R}^N \mid R_0 < |y| < m\}$, which contradicts the fact that $\|v_n\|$ is bounded.

We may assume that $|v_n| \rightarrow 0$ strongly in $L_{\text{loc}}^p(\mathbb{R}^N)$ with $p < 2N/(N-1)$ and thus $|\gamma(v_n)| \rightarrow 0$ strongly in $L_{\text{loc}}^p(\mathbb{R}^N)$.

Let $\xi_\varepsilon \in C^\infty(\mathbb{R}_+^{N+1})$ be a symmetric function, namely $\xi_\varepsilon(x, gy) = \xi_\varepsilon(x, y)$ for all $g \in O(N)$, $x > 0$, $y \in \mathbb{R}^N$. Moreover assume that $\xi_\varepsilon(z) = 0$ if $|z| \leq R_\varepsilon$ and $\xi_\varepsilon(z) = 1$ if $|z| \geq R_\varepsilon + 1$ and $\xi(z) \in [0, 1]$ for all $z \in \mathbb{R}_+^{N+1}$. Set $w_n = \xi_\varepsilon v_n$. We apply now Young's inequality (2.7) with $p = q = 2r/(2r-1)$ and $h = W$, $f = |\gamma(v_n)|^\theta$, $g = |\gamma(v_n)|^\theta - |\gamma(w_n)|^\theta$:

$$\begin{aligned}
& |\mathbb{D}(v_n) - \mathbb{D}(w_n)| \\
& \leq \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) \left| |\gamma(v_n)(x)|^\theta |\gamma(v_n)(y)|^\theta - |\gamma(w_n)(x)|^\theta |\gamma(w_n)(y)|^\theta \right| dx dy \\
& = \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) \left| |\gamma(v_n)(x)|^\theta |\gamma(v_n)(y)|^\theta - |\gamma(v_n)(x)|^\theta |\gamma(w_n)(y)|^\theta \right. \\
& \quad \left. + |\gamma(v_n)(x)|^\theta |\gamma(w_n)(y)|^\theta - |\gamma(w_n)(x)|^\theta |\gamma(w_n)(y)|^\theta \right| dx dy \\
& \leq \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) \left| |\gamma(v_n)(x)|^\theta |\gamma(v_n)(y)|^\theta - |\gamma(w_n)(y)|^\theta \right| dx dy \\
& \quad + \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) \left| |\gamma(w_n)(y)|^\theta |\gamma(v_n)(x)|^\theta - |\gamma(w_n)(x)|^\theta \right| dx dy \\
& \leq 2 \iint_{\mathbb{R}^N \times \mathbb{R}^N} W(x-y) \left| |\gamma(v_n)(x)|^\theta |\gamma(v_n)(y)|^\theta - |\gamma(w_n)(y)|^\theta \right| dx dy \\
& \leq 2C|W|_r |\gamma(v_n)|_{\frac{2r\theta}{2r-1}}^\theta \left| |\gamma(v_n)|^\theta - |\gamma(w_n)|^\theta \right|_{\frac{2r}{2r-1}} = o(1), \tag{4.6}
\end{aligned}$$

since $|\gamma(v_n)|^\theta - |\gamma(w_n)|^\theta \rightarrow 0$ strongly in $L_{\text{loc}}^{\frac{2r}{2r-1}}(\mathbb{R}^N)$. Here and in the following C denotes some positive constant independent of n , not necessarily the same one. Similarly,

$$\begin{aligned}
& \left| \int_{\mathbb{R}^N} \left(W * |\gamma(v_n)|^\theta \right) |\gamma(v_n)|^{\theta-2} \gamma(v_n) \gamma(w_n) \right. \\
& \quad \left. - \int_{\mathbb{R}^N} \left(W * |\gamma(w_n)|^\theta \right) |\gamma(w_n)|^{\theta-2} \gamma(w_n) \gamma(w_n) \right| \\
& \leq 2C|W|_r |\gamma(v_n)|_{\frac{2r\theta}{2r-1}}^\theta \left| |\gamma(v_n)|^\theta - |\gamma(w_n)|^\theta \right|_{\frac{2r}{2r-1}} = o(1).
\end{aligned}$$

Therefore,

$$|I'(v_n)w_n - I'(w_n)w_n| \leq C \iint_{S_\varepsilon} (|\nabla v_n|^2 + m^2 v_n^2) dx dy + \int_{A_\varepsilon} V(y) \gamma(v_n)^2 dy + o(1).$$

Set $u_n = (1 - \xi)v_n$. Analogously we have

$$|I'(v_n)u_n - I'(u_n)u_n| \leq C \iint_{S_\varepsilon} (|\nabla u_n|^2 + m^2 u_n^2) dx dy + \int_{A_\varepsilon} V(y) \gamma(u_n)^2 dy + o(1).$$

Therefore

$$I'(u_n)u_n = O(\epsilon) + o(1) \quad (4.7)$$

and

$$I'(w_n)w_n = O(\epsilon) + o(1). \quad (4.8)$$

From (4.7), we derive that $I(u_n) = \frac{\theta-1}{2\theta}\mathbb{D}(u_n) + O(\epsilon) + o(1) \geq O(\epsilon) + o(1)$.

Consider $t_n > 0$ such that $I'(t_n w_n)(t_n w_n) = 0$ for any n , namely

$$t_n^{2(\theta-1)} = \frac{\|w_n\|^2 + \int_{\mathbb{R}^N} V(y)\gamma(w_n)^2 dy}{\mathbb{D}(w_n)}.$$

From (4.8), we have that $t_n = 1 + O(\epsilon) + o(1)$. Therefore from the characterization of E_α we have

$$\begin{aligned} c + o(1) &= I(v_n) = I(u_n) + I(w_n) + O(\epsilon) \geq I(w_n) + O(\epsilon) + o(1) \\ &\geq I(t_n w_n) + O(\epsilon) + o(1) \geq E_\alpha + O(\epsilon) + o(1). \end{aligned}$$

As $n \rightarrow +\infty$, $\epsilon \rightarrow 0$, we derive that $c \geq E_\alpha$ which is a contradiction. Hence, $c = 0$ and $v_n \rightarrow 0$ strongly in $H^1(\mathbb{R}_+^{N+1})$. $\square$

Lemma 4.2. *Let $\{v_n\}_n$ be a sequence in $H^1(\mathbb{R}_+^{N+1})$ such that $v_n \rightharpoonup v$ weakly in $H^1(\mathbb{R}_+^{N+1})$. The following hold:*

(i) $\mathbb{D}'(v_n)u \rightarrow \mathbb{D}'(v)u$ for all $u \in H^1(\mathbb{R}_+^{N+1})$.

(ii) *After passing to a subsequence, there exists a sequence $\{\tilde{v}_n\}_n$ in $H^1(\mathbb{R}_+^{N+1})$ such that $\tilde{v}_n \rightarrow v$ strongly in $H^1(\mathbb{R}_+^{N+1})$,*

$$\begin{aligned} \mathbb{D}(v_n) - \mathbb{D}(v_n - \tilde{v}_n) &\rightarrow \mathbb{D}(v) \quad \text{in } \mathbb{R}, \\ \mathbb{D}'(v_n) - \mathbb{D}'(v_n - \tilde{v}_n) &\rightarrow \mathbb{D}'(v) \quad \text{in } H^{-1}(\mathbb{R}_+^{N+1}). \end{aligned}$$

Proof. The proof is completely analogous to that of Lemma 3.5 in [1]. The function $\tilde{v}_n$ is the product of v_n with a smooth cut-off function, so $\tilde{v}_n$ belongs to $H^1(\mathbb{R}_+^{N+1})$ if v_n does. We omit the details. $\square$

Proposition 4.3. *The functional $I: H^1(\mathbb{R}_+^{N+1}) \rightarrow \mathbb{R}$ satisfies the Palais-Smale condition $(PS)_c$ at each $c < E_{V_\infty}$, where $V_\infty := \liminf_{|x| \rightarrow \infty} V(x)$.*

Proof. Let $v_n \in H^1(\mathbb{R}_+^{N+1})$ satisfy

$$I(v_n) \rightarrow c < E_{V_\infty} \quad \text{and} \quad I'(v_n) \rightarrow 0$$

strongly in the dual space $H^{-1}(\mathbb{R}_+^{N+1})$. Since $\{v_n\}_n$ is bounded in $H^1(\mathbb{R}_+^{N+1})$ it contains a subsequence such that $v_n \rightharpoonup v$ weakly in $H^1(\mathbb{R}_+^{N+1})$ and $\gamma(v_n) \rightharpoonup \gamma(v)$ in $L^p(\mathbb{R}^N)$ for any $p \in [2, 2N/(N-1)]$.

By Lemma 4.2, v solves (1.3) and, after passing to a subsequence, there exists a sequence $\{\tilde{v}_n\}_n$ in $H^1(\mathbb{R}_+^{N+1})$ such that $u_n := v_n - \tilde{v}_n \rightarrow 0$ weakly in $H^1(\mathbb{R}_+^{N+1})$,

$$\begin{aligned} I(v_n) - I(u_n) &\rightarrow I(v) \quad \text{in } \mathbb{R}, \\ I'(v_n) - I'(u_n) &\rightarrow 0 \quad \text{strongly in } H^{-1}(\mathbb{R}_+^{N+1}). \end{aligned}$$

Hence, $I(v) = \frac{\theta-2}{2\theta} \mathbb{D}(v) \geq 0$,

$$I(u_n) \rightarrow c - I(v) \leq c, \quad \text{and} \quad I'(u_n) \rightarrow 0$$

strongly in $H^{-1}(\mathbb{R}_+^{N+1})$. By Lemma 4.1 a subsequence of $\{u_n\}_n$ converges strongly to 0 in $H^1(\mathbb{R}_+^{N+1})$. This implies that a subsequence of $\{v_n\}_n$ converges strongly to v in $H^1(\mathbb{R}_+^{N+1})$. $\square$

5. Mountain Pass Geometry

Let us consider the limit problem

$$\begin{cases} -\Delta v + m^2 v = 0 & \text{in } \mathbb{R}_+^{N+1} \\ -\frac{\partial v}{\partial x} = -V_\infty v + (W * |v|^\theta) |v|^{\theta-2} v & \text{in } \mathbb{R}^N = \partial \mathbb{R}_+^{N+1} \end{cases} \quad (5.1)$$

where $V_\infty := \liminf_{|x| \rightarrow \infty} V(x) > 0$. By Theorem 3.2, the first mountain pass value E_{V_∞} of the functional J_{V_∞} associated to problem (5.1) is attained at a positive function $\omega_\infty \in H^1(\mathbb{R}_+^{N+1})$, which is symmetric $\omega_\infty(x, gy) = \omega_\infty(x, y)$ for all $g \in O(N)$, $x > 0$, $y \in \mathbb{R}^N$. Moreover, since $V_\infty > 0$, we are allowed to choose $\sigma = 0$, and there exists $C > 0$ such that

$$0 < \omega_\infty(x, y) \leq C e^{-m\sqrt{x^2 + |y|^2}} \quad (5.2)$$

for all $(x, y) \in [0, +\infty) \times \mathbb{R}^N$. In particular, $\gamma(\omega_\infty)$ is radially symmetric in $\mathbb{R}^N$ and

$$0 < \gamma(\omega_\infty)(y) \leq C e^{-m|y|}$$

for any $y \in \mathbb{R}^N$. As in Theorem 3.2, a bootstrap procedure shows that $\omega_\infty \in C^\infty([0, +\infty) \times \mathbb{R}^N)$.

Lemma 5.1. *We have*

$$|\nabla \omega_\infty(z)| = O(e^{-m|z|}) \quad \text{as } |z| \rightarrow \infty. \quad (5.3)$$

Proof. We consider the equation

$$\sqrt{-\Delta + m^2} u + V_\infty u = (W * |u|^\theta) |u|^{\theta-2} u \quad \text{in } \mathbb{R}^N$$

satisfied by ω_∞ . For any index $i = 1, 2, \dots, N$ we write $v_i = \partial \omega_\infty / \partial y_i$ and observe that v_i satisfies

$$\sqrt{-\Delta + m^2} v_i + V_\infty v_i = \theta (W * \omega_\infty^{\theta-1} v_i) \omega_\infty^{\theta-1} + (\theta - 1) (W * \omega_\infty^\theta) \omega_\infty^{\theta-2} v_i \quad (5.4)$$

or, equivalently,

$$\begin{aligned} -\Delta v_i + m^2 v_i &= 0 \quad \text{in } \mathbb{R}_+^{N+1} \\ -\frac{\partial v_i}{\partial x} &= -V_\infty v_i + \theta(W * \omega_\infty^{\theta-1} v_i) \omega_\infty^{\theta-1} + (\theta-1)(W * \omega_\infty^\theta) \omega_\infty^{\theta-2} v_i \quad \text{in } \mathbb{R}^N. \end{aligned}$$

The differentiation of the equation is allowed by the regularity of the solution ω_∞ , see [11, Theorem 3.14]. Moreover, $\omega_\infty \in L^p(\mathbb{R}_+^{N+1})$ for any $p > 1$, because it is bounded and decays exponentially fast at infinity. By elliptic regularity, $\omega_\infty \in W^{2,p}(\mathbb{R}_+^{N+1})$ for any $p > 1$, and in particular $v_i \in L^p(\mathbb{R}_+^{N+1})$ for any $p > 1$. An interpolation estimate shows that $\omega_\infty^{\theta-1} v_i \in L^p(\mathbb{R}_+^{N+1})$ for any $p > 1$. Then the convolution $W * (\omega_\infty^{\theta-1} v_i) \in L^\infty(\mathbb{R}_+^{N+1})$, and the term $(W * (\omega_\infty^{\theta-1} v_i)) \omega_\infty^{\theta-1} \in L^2(\mathbb{R}_+^{N+1})$ by the summability properties of ω_∞ . The term $(W * \omega_\infty^\theta) \omega_\infty^{\theta-2} v_i \in L^2(\mathbb{R}_+^{N+1})$ trivially.

Now the proof of [11, Theorem 3.14] shows that $v_i(x, y) \rightarrow 0$ as $x + |y| \rightarrow +\infty$. A comparison with the function $e^{-m\sqrt{x^2+|y|^2}}$ as in [11, Theorem 5.1] shows the validity of (5.3). $\square$

Fix $\varepsilon \in (0, \frac{2m-k}{2m+k})$. For $R > 0$, we consider a symmetric cut off function $\xi_R \in C^\infty(\mathbb{R}_+^{N+1})$, namely $\xi_R(x, gy) = \xi_R(x, y)$ for all $g \in O(N)$, $x > 0$, $y \in \mathbb{R}^N$ such that $\xi_R(z) = 0$ if $|z| \geq R$ and $\xi_R(z) = 1$ if $|z| \leq R(1-\varepsilon)$ and $\xi_R(z) \in [0, 1]$ for all $z \in \mathbb{R}_+^{N+1}$.

Let us define $\omega^R(z) := \omega_\infty(z) \xi_R(z)$, for any $z \in \mathbb{R}_+^{N+1}$.

Lemma 5.2. *As $R \rightarrow \infty$,*

$$\left| \iint_{\mathbb{R}_+^{N+1}} |\nabla \omega_\infty|^2 - |\nabla \omega^R|^2 \right| = O(R^{N-1} e^{-2m(1-\varepsilon)R}), \quad (5.5)$$

$$|\mathbb{D}(\omega_\infty) - \mathbb{D}(\omega^R)| = O(R^{N-1} e^{-\theta m(1-\varepsilon)R}). \quad (5.6)$$

Proof. The proof of (5.5) is standard. Indeed, using (5.3) and cylindrical coordinates in $\mathbb{R}_+^{N+1}$,

$$\begin{aligned} \left| \iint_{\mathbb{R}_+^{N+1}} |\nabla \omega^R|^2 - |\nabla \omega_\infty|^2 \right| &\leq C \iint_{\{z \in \mathbb{R}_+^{N+1} \mid (1-\varepsilon)R < |z| \leq R\}} |\nabla \omega_\infty|^2 \\ &\leq C_1 \iint_{\{z \in \mathbb{R}_+^{N+1} \mid (1-\varepsilon)R < |z| \leq R\}} e^{-2m|z|} dz \\ &\leq C_1 R^{N-1} e^{-2m(1-\varepsilon)R}. \end{aligned}$$

To prove (5.6), we recall that $W = W_1 + W_2 \in L^r(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$. The difference $\mathbb{D}(\gamma(\omega_\infty)) - \mathbb{D}(\gamma(\omega^R))$ can be split in two parts, the one with W_1 and the one with

W_2 . The former can be estimated as follows:

$$\begin{aligned}
& |\mathbb{D}(\gamma(\omega_\infty)) - \mathbb{D}(\gamma(\omega^R))| \\
& \leq \int_{\mathbb{R}^N \times \mathbb{R}^N} |\gamma(\omega_\infty)(x)|^\theta |\gamma(\omega_\infty)(y)|^\theta - |\gamma(\omega^R)(x)|^\theta |\gamma(\omega^R)(y)|^\theta |W_1(x-y) dx dy \\
& \leq 2 \int_{\mathbb{R}^N \times \mathbb{R}^N} W_1(x-y) |\gamma(\omega_\infty)(x)|^\theta ||\gamma(\omega_\infty)(y)|^\theta - |\gamma(\omega^R)(y)|^\theta| dx dy \\
& \leq 2 ||\gamma(\omega_\infty)|^\theta - |\gamma(\omega^R)|^\theta||_{\frac{2r}{2r-1}} \|\gamma(\omega_\infty)\|_{\frac{2r\theta}{2r-1}}^\theta \|W_1\|_r \\
& \leq C \left(\int_{(1-\varepsilon)R}^\infty t^{N-1} e^{-m\frac{2r\theta}{2r-1}t} dt \right)^{\frac{2r-1}{2r}} = C_2 R^{N-1} e^{-\theta m(1-\varepsilon)R}.
\end{aligned}$$

The latter is simpler, since we use directly the L^∞ -norm of W_2 . $\square$

For $s \in \mathbb{R}^N$, set $R_s := \frac{k+2m}{4m} |s|$. Since $k \in (0, 2m)$, it results that $R_s \in (0, |s|)$. Hence $|s| - R_s \rightarrow +\infty$, as $|s| \rightarrow +\infty$. With this notation, we define the function

$$\omega_s^{R_s}(z) := \omega_\infty(x, y-s) \xi_{R_s}(x, y-s)$$

where $z = (x, y) \in \mathbb{R}^{N+1}$.

Lemma 5.3. *There exist $\varrho_0, d_0 \in (0, \infty)$ such that*

$$I(t(\omega_s^{R_s})) \leq E_{V_\infty} - d_0 e^{-k|y|} \quad \text{for all } t \geq 0,$$

provided that $|s| \geq \varrho_0$

Proof. For $u \in H^1(\mathbb{R}_+^{N+1})$ we have by (2.3) that $\max_{t \geq 0} I(tu) = I(t_u u)$ if and only if

$$t_u = \left(\frac{\|u\|^2 + \int_{\mathbb{R}^N} V(y) \gamma(u)^2 dy}{\mathbb{D}(u)} \right)^{1/(2\theta-2)}.$$

Indeed

$$\|u\|^2 + \int_{\mathbb{R}^N} V(y) \gamma(u)^2 dy \geq \left(1 - \frac{V_0}{m}\right) \int_{\mathbb{R}_+^{N+1}} |\nabla u|^2 + m(m - V_0) \int_{\mathbb{R}_+^{N+1}} |u|^2 > 0. \quad (5.7)$$

So, since $\omega_\infty^{R_s} \rightarrow \omega_\infty$ in $H^1(\mathbb{R}_+^{N+1})$ as $|s| \rightarrow \infty$, and taking into account that $I_{V_\infty}(\omega_\infty) = \max_{t \geq 0} I_{V_\infty}(t(\omega_\infty))$ there exist $0 < t_1 < t_2 < +\infty$ such that

$$\max_{t \geq 0} I(t(\omega_s^{R_s})) = \max_{t_1 \leq t \leq t_2} I(t(\omega_s^{R_s}))$$

for all large enough $|s|$.

Let $t \in [t_1, t_2]$. Write $V = V^+ - V^-$, where $V^+(x) = \max\{V(x), 0\}$ and $V^-(x) = \max\{-V(x), 0\}$, and remark that the assumption $V_\infty > 0$ implies $V(x) = V^+(x)$

whenever $|x|$ is sufficiently large. Assumption (V_2) yields therefore

$$\begin{aligned} \int_{\mathbb{R}^N} V(y) (t\gamma(\omega_s^{R_s}))^2(y) dy &\leq t^2 \int_{|y| \leq R_s} V^+(y+s) (\gamma(\omega_s^{R_s}))^2(y) dy \\ &\leq t^2 \int_{|y| \leq R_s} (V_\infty - c_0 e^{-k|y+s|}) (\gamma(\omega_\infty))^2(y) dy \\ &\leq \int_{\mathbb{R}^N} V_\infty (t\gamma(\omega_\infty))^2 - \left(c_0 t_1^2 \int_{|y| \leq 1} e^{-k|y|} (\gamma(\omega_\infty))^2(y) dy \right) e^{-k|s|} \end{aligned}$$

for $|s|$ large enough.

Therefore, using Lemma 5.2, we get

$$\begin{aligned} I(t(\omega_s^{R_s})_s) &= \frac{1}{2} \|t(\omega_s^{R_s})_s\|^2 + \frac{1}{2} \int_{\mathbb{R}^N} V(y) (t\gamma(\omega_\infty))^2 dy - \frac{1}{2\theta} \mathbb{D}(t\omega_s^{R_s}) \\ &\leq \frac{1}{2} \|t\omega_\infty\|^2 + \frac{1}{2} \int_{\mathbb{R}^N} V_\infty (t\gamma(\omega_\infty))^2 dy - \frac{1}{2\theta} \mathbb{D}(t\omega_\infty) \\ &\quad - C e^{-k|s|} + O(R_s^{N-1} e^{-2m(1-\varepsilon)R_s}) \\ &\leq \max_{t \geq 0} I_{V_\infty}(t\omega_\infty) - d_0 e^{-\kappa|s|} \\ &= E_{V_\infty} - d_0 e^{-k|s|} \end{aligned}$$

for sufficiently large $|s|$, because our choices of ε and R_s guarantee that $2m(1-\varepsilon)R_s > k|s|$. $\square$

6. Proof of theorem 1.1

The proof of Theorem 1.1 is now immediate. The Euler functional I satisfies the geometric assumptions of the Mountain Pass Theorem (see [2]) on $H^1(\mathbb{R}_+^{N+1})$. Since it also satisfies the Palais-Smale condition, as we showed in the previous sections, we conclude that I possesses at least a critical point $v \in H^1(\mathbb{R}_+^{N+1})$. In addition,

$$I(v) = c = \inf_{\gamma \in \Gamma} \max_{0 \leq t \leq 1} I(\gamma(t)),$$

where $\Gamma = \{\gamma \in C([0, 1], H^1(\mathbb{R}_+^{N+1})) \mid \gamma(0) < 0, I(\gamma(1)) < 0\}$.

To prove that $v \geq 0$, we notice that, reasoning as in (5.7), the map $t \mapsto I(tw)$ has one and only one strict maximum point at $t = 1$ whenever $w \in H^1(\mathbb{R}_+^{N+1})$ is a critical point of I . Since $I(|w|) \leq I(w)$ for all $w \in H^1(\mathbb{R}_+^{N+1})$, and

$$I(t|w|) \leq I(tw) < I(w) \quad \text{for every } t > 0, t \neq 1,$$

we conclude that

$$c \leq \sup_{t \geq 0} I(t|v|) \leq I(v) = c.$$

We claim that $|v|$ is also a critical point of I . Indeed, otherwise, we could deform the path $t \mapsto t|v|$ into a path $\gamma \in \Gamma$ such that $I(\gamma(t)) < c$ for every $t \geq 0$, a contradiction with the definition of c .

7. Further properties of the solution

We collect in the next statement some additional features of the weak solution found above.

Theorem 7.1. *Let u be the solution to equation (1.3) provided by Theorem 1.1. Then $u \in C^\infty(\mathbb{R}^N) \cap L^q(\mathbb{R}^N)$ for every $q \geq 2$. Moreover,*

$$0 < u(y) \leq Ce^{-m|y|}. \quad (7.1)$$

Proof. The regularity of u can be established by mimicking the proofs in Section 3 of [11]. The potential function V is harmless, being bounded from above and below.

To prove the exponential decay at infinity, we introduce a comparison function

$$W_R(x, y) = C_R e^{-m\sqrt{x^2 + |y|^2}}, \quad \text{for every } (x, y) \in \mathbb{R}_+^{N+1},$$

and we will fix $R > 0$ and $C_R > 0$ in a suitable manner. We also introduce the notation

$$\begin{aligned} B_R^+ &= \left\{ (x, y) \in \mathbb{R}_+^{N+1} \mid \sqrt{x^2 + |y|^2} < R \right\} \\ \Omega_R^+ &= \left\{ (x, y) \in \mathbb{R}_+^{N+1} \mid \sqrt{x^2 + |y|^2} > R \right\} \\ \Gamma_R &= \left\{ (0, y) \in \partial\mathbb{R}_+^{N+1} \mid |y| \geq R \right\}. \end{aligned}$$

It is easily seen that

$$\begin{cases} -\Delta W_R + m^2 W_R \geq 0 & \text{in } \Omega_R^+ \\ -\frac{\partial W_R}{\partial x} = 0 & \text{on } \Gamma_R^+. \end{cases}$$

Call $w(x, y) = W_R(x, y) - v(x, y)$, and remark that $-\Delta w + m^2 w \geq 0$ in Ω_R^+ . If $C_R = e^{mR} \max_{\partial B_R^+} v$, then $w \geq 0$ on ∂B_R^+ and $\lim_{x+|y| \rightarrow +\infty} w(x, y) = 0$. We claim that $w \geq 0$ in the closure $\overline{\Omega_R^+}$.

If not, $\inf_{\Omega_R^+} w < 0$, and the strong maximum principle provides a point $(0, y_0) \in \Gamma_R$ such that

$$w(0, y_0) = \inf_{\Omega_R^+} w < w(x, y) \quad \text{for every } (x, y) \in \Omega_R^+.$$

For some $0 < \lambda < m$, we introduce $z(x, y) = w(x, y)e^{\lambda x}$. As before, $\lim_{x+|y| \rightarrow +\infty} z(x, y) = 0$ and $z \geq 0$ on ∂B_R^+ . Since

$$0 \leq -\Delta w + m^2 w = e^{-\lambda x} \left(-\Delta z + 2\lambda \frac{\partial z}{\partial x} + (m^2 - \lambda^2)z, \right)$$

the strong maximum principle applies and yields that $\inf_{\Gamma_R} z = \inf_{\Omega_R^+} z < z(x, y)$ for every $(x, y) \in \Omega_R^+$. Therefore $z(0, y_0) = \inf_{\Gamma_R} z = \inf_{\Gamma_R} w < 0$. Hopf's lemma now gives

$$-\frac{\partial w}{\partial x}(0, y_0) - \lambda w(0, y_0) < 0.$$

But this is impossible. Indeed

$$-\frac{\partial w}{\partial x}(0, y_0) = -V(y_0)v(0, y_0) - (W * |v|^\theta)|v(0, y_0)|^{\theta-2}v(0, y_0),$$

and hence

$$-\frac{\partial w}{\partial x}(0, y_0) - \lambda v(0, y_0) = -\lambda v(0, y_0) - V(y_0)v(0, y_0) - (W * |v|^\theta)|v(0, y_0)|^{\theta-2}v(0, y_0).$$

Recall that $v(0, y_0) < 0$ and $\lambda > 0$; if we can show that

$$-V(y_0)v(0, y_0) - (W * |v|^\theta)|v(0, y_0)|^{\theta-2}v(0, y_0) \geq 0,$$

we will be done. First of all, we recall that (see [11, pag. 70] and also [9, Lemma 2.3])

$$\lim_{|y| \rightarrow +\infty} (W * |v|^\theta)|v(0, y)|^{\theta-2}v(0, y) = 0,$$

since $\lim_{|y| \rightarrow +\infty} W(y) = 0$. So we pick $R > 0$ so large that $|(W * |v|^\theta)|v(0, y_0)|^{\theta-2}v(0, y_0)|$ is very small. Choosing R even larger, we can also assume that $V(y_0) > 0$, since $V_\infty > 0$. Hence $-V(y_0)v(0, y_0) - (W * |v|^\theta)|v(0, y_0)|^{\theta-2}v(0, y_0) \geq 0$, and the proof is finished.

To summarize, we have proved that, whenever $x + |y|$ is sufficiently large, then

$$v(x, y) \leq W_R(x, y),$$

and hence the validity of (7.1).

Acknowledgements. The first author is supported by MIUR Project (PRIN 2009) “Variational and topological methods in the study of nonlinear phenomena” and by GNAMPA (INDAM) Project 2013 “Problemi differenziali di tipo ellittico nei fenomeni fisici non lineari”. The second author is supported by MIUR Project (PRIN 2009) “Teoria dei punti critici e metodi perturbativi per equazioni differenziali nonlineari”.

□

References

- 1 N. Ackermann, *On a periodic Schrödinger equation with nonlocal superlinear part*, Math. Z. **248** (2004), 423–443.
- 2 A. Ambrosetti, P.H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973), 349–381.
- 3 X. Cabré, J. Solà-Morales, *Layers solutions in a half-space for boundary reactions*, Comm. Pure Applied Math. **58** (2005), 1678–1732.
- 4 X. Cabré, J. Tan, *Positive solutions of nonlinear problems involving the square root of the Laplacian*, Adv. Math. **224** (2010), 2052–2093.
- 5 Y. Cho, T. Ozawa, *On the semirelativistic Hartree-type equation*, SIAM J. Math. Anal. **38** (2006), no.4, 1060–1074.
- 6 L. Caffarelli, L. Silvestre, *An extension problem related to the fractional Laplacian*, Communications in Partial Differential Equations **32** (2007), 1245–1260.

- 7 S. Cingolani, M. Clapp, S. Secchi, *Multiple solutions to a magnetic nonlinear Choquard equation*, Zeitschrift für Angewandte Mathematik und Physik (ZAMP), **63** (2012), 233–248.
- 8 S. Cingolani, M. Clapp, S. Secchi, *Intertwining semiclassical solutions to a Schrödinger-Newton system*, Discrete Continuous Dynamical Systems Series S **6** (2013), 891–908.
- 9 S. Cingolani, S. Secchi, M. Squassina, *Semiclassical limit for Schrödinger equations with magnetic field and Hartree-type nonlinearities*, Proc. Roy. Soc. Edinburgh **140 A** (2010), 973–1009.
- 10 V. Coti Zelati, M. Nolasco, *Ground states for pseudo-relativistic Hartree equations of critical type*, preprint 2012. <http://arxiv.org/abs/1202.2446>
- 11 V. Coti Zelati, M. Nolasco, *Existence of ground state for nonlinear, pseudorelativistic Schrödinger equations*, Rend. Lincei Mat. Appl. **22** (2011), 51–72.
- 12 A. Elgart, B. Schlein, *Mean field dynamics of boson stars*, Comm. Pure Appl. Math. **60** (2007), 500–545.
- 13 J. Fröhlich, E. Lenzmann, *Mean-field limit of quantum Bose gases and nonlinear Hartree equation*, in Séminaire: Équations aux Dérivées Partielles 2003–2004, Exp. No. XIX, 26 pp., École Polytech., Palaiseau, 2004.
- 14 J. Fröhlich, J. Jonsson, E. Lenzmann, *Boson stars as solitary waves*, Comm. Math. Phys. **274** (2007), 1–30.
- 15 E. Lenzmann, *Well-posedness for Semi-relativistic Hartree equations with Critical type*, Math. Phys. Anal. Geom. **10** (2007), 43–64.
- 16 E. Lenzmann, *Uniqueness of ground states for pseudo relativistic Hartree equations*, Analysis and PDE **2** (2009), 1–27.
- 17 E.H. Lieb, *Existence and uniqueness of the minimizing solution of Choquard’s nonlinear equation*, Stud. Appl. Math. **57** (1977), 93–105.
- 18 E.H. Lieb, B. Simon, *The Hartree-Fock theory for Coulomb systems*, Comm. Math. Phys. **53** (1977), 185–194.
- 19 E.H. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, The Annals of Mathematics **118** (1983), 349–374.
- 20 E.H. Lieb, H.-T. Yau, *The Chandrasekhar theory of stellar collapse as the limit of quantum mechanics*, Comm. Math. Phys. **112** (1987), 147–174.
- 21 P.-L. Lions, *The Choquard equation and related questions*, Nonlinear Anal. T.M.A. **4** (1980), 1063–1073.
- 22 L. Ma, L. Zhao, *Classification of positive solitary solutions of the nonlinear Choquard equation*, Arch. Rational Mech. Anal. **195** (2010), 455–467.
- 23 M. Melgaard, F. Zongo, *Multiple solutions of the quasirelativistic Choquard equation*, J. Math. Phys. **53** (2012), no. 3, 033709, 12 pp.
- 24 I.M. Moroz, R. Penrose, P. Tod, *Spherically-symmetric solutions of the Schrödinger-Newton equations*, Topology of the Universe Conference (Cleveland, OH, 1997), Classical Quantum Gravity **15** (1998), 2733–2742.
- 25 I.M. Moroz, P. Tod, *An analytical approach to the Schrödinger-Newton equations*, Nonlinearity **12** (1999), 201–216.
- 26 D. Mugnai, *The pseudorelativistic Hartree equation with a general nonlinearity: existence, non existence and variational identities*, preprint <http://arxiv.org/abs/1206.0853>.
- 27 R. Penrose, *Quantum computation, entanglement and state reduction*, R. Soc. Lond. Philos. Trans. Ser. A Math. Phys. Eng. Sci. **356** (1998), 1927–1939.
- 28 R. Penrose, *The road to reality. A complete guide to the laws of the universe*, Alfred A. Knopf Inc., New York 2005.
- 29 S. Secchi, *A note on Schrödinger-Newton systems with decaying electric potential*, Nonlinear Analysis **72** (2010), 3842–3856.
- 30 L. Tartar, *An introduction to Sobolev spaces*. Springer, 2007.
- 31 P. Tod, *The ground state energy of the Schrödinger-Newton equation*, Physics Letters A **280** (2001), 173–176.
- 32 J. Wei, M. Winter, *Strongly interacting bumps for the Schrödinger-Newton equation*, J. Math. Phys. **50** (2009), 012905.
- 33 M. Willem, *Minimax theorems. Progress in Nonlinear Differential Equations and their Applications*, 24. Birkhäuser Boston, Inc., Boston, MA, 1996.